



CASE REPORT

Open Access

# Anteroposterior chest radiograph vs. chest CT scan in early detection of pneumothorax in trauma patients

Hesham R Omar<sup>1,3\*</sup>, Devanand Mangar<sup>2,3</sup>, Suneel Khetarpal<sup>4</sup>, David H Shapiro<sup>4</sup>, Jaya Kolla<sup>3</sup>, Rania Rashad<sup>5</sup>, Engy Helal<sup>6</sup> and Enrico M Camporesi<sup>3,7</sup>

## Abstract

Pneumothorax is a common complication following blunt chest wall trauma. In these patients, because of the restrictions regarding immobilization of the cervical spine, Anteroposterior (AP) chest radiograph is usually the most feasible initial study which is not as sensitive as the erect chest X-ray or CT chest for detection of a pneumothorax. We will present 3 case reports which serve for better understanding of the entity of occult pneumothorax. The first case is an example of a true occult pneumothorax where an initial AP chest X-ray revealed no evidence of pneumothorax and a CT chest immediately performed revealed evidence of pneumothorax. The second case represents an example of a missed rather than a truly occult pneumothorax where the initial chest radiograph revealed clues suggesting the presence of pneumothorax which were missed by the reading radiologist. The third case emphasizes the fact that "occult pneumothorax is predictable". The presence of subcutaneous emphysema and pulmonary contusion should call for further imaging with CT chest to rule out pneumothorax. Thoracic CT scan is therefore the "gold standard" for early detection of a pneumothorax in trauma patients. This report aims to sensitize readers to the entity of occult pneumothorax and create awareness among intensivists and ER physicians regarding the proper diagnosis and management.

## Introduction

The concept of occult pneumothorax is well accepted among the surgical trauma literature [1-5]. In trauma patients, because of restrictions regarding cervical spine immobilization, AP chest radiograph is usually utilized to detect intrathoracic pathology. This report emphasizes how AP chest radiograph can dangerously delay the recognition of a pneumothorax. More advanced imaging modalities including Chest CT scan or ultrasonography is therefore mandatory to exclude the diagnosis.

## Case # 1

A 24-year-old male presented to the ER after a motor vehicle accident. On admission the patient was confused with a Glasgow coma score of 14/15. CT brain revealed

brain edema and fracture skull base. Chest exam and arterial blood gases were satisfactory. AP chest X-ray revealed no evidence of pneumothorax as demonstrated in Figure 1 panel A, adapted from Omar et. al. [6] CT chest performed immediately after the X-ray revealed a right sided pneumothorax (Figure 1 panel B), adapted from Omar et. al [6]

This case represents an example of a true occult pneumothorax where an AP chest X-ray failed to show an existing pneumothorax. This emphasizes the importance of chest CT in any trauma victim who is tachypnic or hypoxic when the initial AP chest radiograph appears normal. This is especially important in patients expected to be maintained on positive pressure ventilation.


## Case # 2

A 29-year-old restrained driver was involved in a T-bone vehicular accident. At the scene of the accident the patient's Glasgow coma score was 4/15. The patient

\* Correspondence: hesham\_omar2003@yahoo.com

<sup>1</sup>Departement of Internal Medicine, Mercy Hospital and Medical Center, Chicago, Illinois, USA


Full list of author information is available at the end of the article



**Figure 1 Anteroposterior chest X-ray and CT scan chest demonstrating "true occult pneumothorax".** Anteroposterior chest X-ray revealing no evidence of pneumothorax (Panel A). CT chest performed immediately after X-ray revealing right sided pneumothorax (Panel B). Adapted from Omar et. al. [6].

was intubated for airway protection and sent to the ER. While in the ER, an AP chest X-ray was completed (Figure 2 panel a), adapted from Omar et. al, [6] which revealed a mechanically ventilated patient with diffuse airspace opacities prominently located in the left lower lung field. In the setting of trauma, this was interpreted as lung contusions.

The patient was immediately sent for a chest CT scan (Figure 2 panel b), adapted from Omar et. al, [6] that was performed 15 minutes after the chest X-ray and revealed a large left sided pneumothorax, a left lower lobe collapse as well as hemorrhagic contusions of both lower lobes. The anesthesiology/critical care team then decided to insert a chest tube which was



**Figure 2 Anteroposterior chest X-ray and CT scan chest demonstrating "missed pneumothorax".** Initial Anteroposterior chest X-ray of the intubated patient, illustrating diffuse air space opacities in the left lower lung field. Underlying pneumothorax was suggested because of a visible pleural stripe in the lung apex and a highly visible cardiophrenic sulcus (Panel A). Chest CT scan illustrating a left-sided pneumothorax with underlying lung collapse (Panel B). Adapted from Omar et. al. [6].

followed by relief of the pneumothorax and lung re-expansion.

This case represents an example of a missed rather than a truly occult pneumothorax. The presence of a visible pleural stripe at the lung apex and a highly visible cardiophrenic sulcus should have drawn attention to an underlying pneumothorax. This case emphasizes the importance of interpretation of the X-ray both by an experienced radiologist as well as the managing trauma team to avoid missing a pneumothorax.

### Case # 3


A 42-year-old male presented to the ER after a motor vehicle accident. On admission, his Glasgow coma score was 7/15 and CT brain revealed a left thalamic hemorrhage, subarachnoid hemorrhage and brain edema. Chest examination revealed equal air entry on both sides with coarse ronchi and crepitations over the left hemithorax. There was clinical evidence of subcutaneous emphysema in the left chest wall. The patient was hypoxic on room air and an arterial blood gases revealed a PH of 7.35, PO<sub>2</sub> 66.4 mmHg, PCO<sub>2</sub> 33.4 mmHg, HCO<sub>3</sub> 18.2 and SO<sub>2</sub> 91.4% so the patient was maintained on supplemental oxygen of 10 liters/minute supplied by a face mask. AP chest X-ray revealed pulmonary contusions more on the left lung and left sided subcutaneous emphysema with no clear evidence of pneumothorax (Figure 3 panel a), adapted from Omar et. al. [6] Chest CT scan immediately performed revealed left sided pneumothorax and an intercostal

tube was inserted, (Figure 3 panel b), adapted from Omar et. al, [6]

This case illustrates the predictability of occult pneumothorax. The presence of subcutaneous emphysema and pulmonary contusions highly predicts for the presence of an underlying pneumothorax. We therefore suggest that any trauma victim presenting with subcutaneous emphysema, pulmonary contusion or rib fractures should be further investigated with a CT chest if the chest X-ray did not reveal pneumothorax. This is especially important in patients who will receive mechanical ventilation for fear of development of tension pneumothorax.

### Discussion

The widespread availability and utilization of CT scan in the evaluation of trauma patients accounted for the diagnosis of pneumothoraces that were not initially evident on the chest X-ray. The reported incidence of occult pneumothorax varies between 3.7% in injured children presenting to an emergency department to 64% in intubated multi-trauma patients [7-9]. The proportion of pneumothoraces that are occult compared with those actually seen on the chest X-ray ranges from 29% to 72% [1-4,10,11]. Most instances of occult pneumothoraces in the supine patient are anterior and usually located in the superior thorax or thoracic apex indicating that the gold standard for ruling out occult pneumothorax is a thoracic CT scan. Occult pneumothorax is more concerning in trauma patients who



**Figure 3 Anteroposterior chest X-ray and CT scan chest demonstrating the predictability of occult pneumothorax.** AP chest X-ray revealing evidence of bilateral lung contusions and left subcutaneous emphysema (Panel A). Chest CT confirming both the lung contusions and the subcutaneous emphysema and demonstrating left sided pneumothorax not initially appearing on the anteroposterior chest X-ray (Panel B). Adapted from Omar et. al.[6].

have diminished cardiopulmonary reserve because of the risk of rapid progression to tension pneumothorax especially those on positive pressure ventilation.

The traditional management of post-traumatic pneumothoraces has been the placement of a chest tube [5]. The choice between close observation vs. early intercostal tube placement is still unsettled. In patients who are asymptomatic, not mechanically ventilated, with a pneumothorax that is not increasing in size, clinical opinion supports that close observation is safe [3,4,12,13]. However, some authors believe that the risk of progression to tension pneumothorax is significant and that prophylactic chest tube placement is necessary for any patient subjected to positive-pressure ventilation is necessary [10,14,15]. Of note is that the size of the initial occult pneumothorax does not predict for the development of tension pneumothorax and therefore should not be used as a guide for chest tube placement [16].

The 3 case presentations well illustrated the entity of occult pneumothorax. The first case is a classic example of a true occult pneumothorax which did not show initially on the AP chest X-ray and was diagnosed by chest CT scan. Missing the diagnosis is extremely dangerous for fear of the development of tension pneumothorax especially if the patient is placed on a mechanical ventilator. The second case is not a true occult pneumothorax and is better nominated as a missed pneumothorax where the X-ray report did not confirm the presence of pneumothorax when there were 2 clues for the diagnosis. The presence of an apical pleural stripe and a clearly visible cardiophrenic sulcus should have drawn attention to the diagnosis. We therefore suggest that the X-ray should be interpreted by a well trained radiologist and also by the treating physician who, in the setting of a trauma patient, will have a higher index of suspicion for the diagnosis. The third case demonstrated the predictability of occult pneumothorax. The 3 factors that have a predictive value for occult pneumothorax are the presence of subcutaneous emphysema, lung contusions and rib fractures [17]. In this patient the presence of subcutaneous emphysema directed our attention for further imaging looking for a pneumothorax despite its absence on the AP chest radiograph. It should also be noted that the surgical emphysema can obscure the underlying pleural air further limiting the accuracy of the Chest X-ray in diagnosing pneumothorax in such cases.

We conclude that thoracic CT scan is the "gold standard" for early detection of a pneumothorax in blunt trauma patients. In centers utilizing eFAST (Extended Focused Assessment with Sonography for Trauma) technology for trauma victims, this should be the initially utilized modality because of its high sensitivity, no need

for transporting an unstable patient and its low radio-carcinogenesis profile.

## Consent

Written informed consent was obtained from the patient's relatives for publication of this article. A copy of the written consent is available for review by the Editor-in-Chief of this journal

## Author details

<sup>1</sup>Departement of Internal Medicine, Mercy Hospital and Medical Center, Chicago, Illinois, USA. <sup>2</sup>Department of Anesthesiology; Tampa General Hospital; Tampa, Florida, USA. <sup>3</sup>Florida Gulf to Bay Anesthesiology, Tampa, Florida, USA. <sup>4</sup>Department of Surgery, University of South Florida, Tampa, Florida, USA. <sup>5</sup>Department of Critical Care Medicine, Cairo University Hospital, Cairo, Egypt. <sup>6</sup>Emergency Department, Elagouza Hospital, Cairo, Egypt. <sup>7</sup>Department of Surgery/Anesthesiology, University of South Florida, Tampa, Florida, USA.

## Authors' contributions

HO is responsible for literature search and drafting the manuscript and providing the explanatory figures. JK, EC and DM, RR, DS, SK and EH have made critical revisions to the manuscript. All authors have read and approved the whole manuscript.

## Competing interests

The authors declare that they have no competing interests.

Received: 25 May 2010 Accepted: 27 September 2011

Published: 27 September 2011

## References

1. Neff MA, Monk JS, Peters K, Nikhilesh A: **Detection of occult pneumothoraces on abdominal computed tomographic scans in trauma patients.** *J Trauma* 2000, **49**:281-285.
2. Ball CG, Hameed SM, Evans D, Kortbeek JB, Kirkpatrick AW, Canadian Trauma Trials Collaborative: **Occult pneumothorax in the mechanically ventilated trauma patient.** *Can J Surg* 2003, **46**:373-379.
3. Brasel KJ, Stafford RE, Weigelt JA, Tenquist JE, Borgstrom DC: **Treatment of occult pneumothoraces from blunt trauma.** *J Trauma* 1999, **46**:987-991.
4. Hill SL, Edmisten T, Holtzman G, Wright A: **The occult pneumothorax: an increasing entity in trauma.** *Am Surg* 1999, **65**:254-258.
5. American College of Surgeons Committee on Trauma: **Advanced Trauma Life Support Course for Doctors: Instructors Course Manual.** Chicago: American College of Surgeons; 1997.
6. Omar HR, Abdelmalak H, Mangar D, Rashad R, Helal E, Camporesi EM: **Occult pneumothorax, revisited.** *Journal of Trauma Management & Outcomes* 2010, **4**(1):12.
7. Guerrero-Lopez F, Vasquez-Mata G, Alcazar-Romero P, Fernandez-Mondejar E, Aguayo-Hoyes E, Linde-Valverde CM: **Evaluation of the utility of computed tomography in the initial assessment of the critical care patient with chest trauma.** *Crit Care Med* 2000, **28**:1370-1375.
8. Voggenreiter G, Aufmkolk M, Majetschak M, Assenmacher S, Waydhas C, Obertacke U, Nast-Kolb D: **Efficacy of chest computed tomography in critically ill patients with multiple trauma.** *Crit Care Med* 2000, **28**:1033-1039.
9. Kirkpatrick AW, Sirois M, Laupland KB, Liu D, Rowan K, Ball CG, Hameed SM, Brown R, Simons R, Dulchavsky SA, Hamilton DR, Nicolaou S: **Hand-held thoracic sonography for detecting post-traumatic pneumothoraces: The extended focused assessment with sonography for trauma (eFAST).** *J Trauma* 2004, **57**:288-295.
10. Trupka A, Waydhas C, Hallfeldt KJ, Nast-Kolb D, Pfeifer KJ, Schweiberer L: **Value of thoracic computed tomography in the first assessment of severely injured patients with blunt chest trauma: Results of a prospective study.** *J Trauma* 1997, **43**:405-412.
11. Holmes JF, Brant WE, Bogren HG, London KL, Kuppermann N: **Prevalence and importance of pneumothoraces visualized on abdominal computed**

- tomographic scan in children with blunt trauma. *J Trauma* 2001, **50**:516-520.
12. Collins JC, Levine G, Waxman K: **Occult traumatic pneumothorax: immediate tube thoracostomy versus expectant management.** *Am Surg* 1992, **58**:743-746.
  13. Wolfman NT, Myers MS, Glauser SJ, Meredith JW, Chen MY: **Validity of CT classification on management of occult pneumothorax: A prospective study.** *AJR Am J Roentgenol* 1998, **171**:1317-1323.
  14. Omert L, Yeane WW, Protech J: **Efficacy of thoracic computerized tomography in blunt chest trauma.** *Am Surg* 2001, **67**:660-667.
  15. Karnik AM, Khan FA: **Pneumothorax and barotrauma.** In *Critical Care Medicine: Principles of Diagnosis and Management in the Adult*. 2 edition. Edited by: Parillo JE, Dellinger RP. St. Louis: Mosby; 2001:930-948.
  16. Enderson BL, Abdalla R, Frame SB, Casey MT, Gould H, Maull KI: **Tube thoracostomy for occult pneumothorax: a prospective randomized study of its use.** *J Trauma* 1993, **35**:726-730.
  17. Ball CG, Kirkpatrick AW, Laupland KB, *et al*: **Incidence, risk factors and outcomes for occult pneumothoraces in victims of major trauma.** *J Trauma* 2005, **59**:917-25.

doi:10.1186/1755-7682-4-30

**Cite this article as:** Omar *et al.*: Anteroposterior chest radiograph vs. chest CT scan in early detection of pneumothorax in trauma patients. *International Archives of Medicine* 2011 **4**:30.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

