



ORIGINAL RESEARCH

Open Access

Profile of idiopathic parkinson's disease in Moroccan patients

Wafa Regragui^{1,5†}, Lamiae Lachhab^{1,5*†}, Rachid Razine^{2,4,5}, Leila Raissouni^{1,5}, Khaoula Rasmouni^{1,5}, Fatima Imounan^{1,5}, El hachmia Ait Benhaddou^{1,5}, Redouane Abouqal^{2,5}, Ali Benomar^{1,3,5} and Mohamed Yahyaoui^{1,5}

Abstract

Objective: To characterize clinical aspects of Idiopathic Parkinson's disease from a movement disorders consultation in University Hospital of Rabat.

Methods: Retrospective review of medical records of 117 patients with diagnosis of Idiopathic Parkinson's disease seen in our outpatient clinic from 2006 to 2011.

Results: Mean age was 64 ± 10 years with predominance of men (61.5%). Mean age at disease onset was 57 ± 11 years. Early onset Parkinson's Disease was recorded in 12.8%. The median duration of disease was 5 years. Initial symptom appeared on the right side in 56.5%. Tremor presentation was the most frequent (40.2%). Symptom severity was mild to moderate in 80% of cases (UPDRS < 30). Forty four per cent of patients were receiving both Dopamine Agonists and Levodopa and in 69% of cases Levodopa was introduced within the first year following onset. The mean Levodopa Equivalent Doses (LED) was 667 ± 446 mg/day. Motor complications were found in 42% with motor fluctuations in 28.7% and ²dyskinesias in 26.7%. Non motor complications are represented mainly by autonomic disorders (44%). There were no differences in the clinical presentation related to the age at onset. Age of onset < 45 and LED > 600 mg are identified as risk factors for motor fluctuations whereas duration of Levodopa treatment is a risk factor of dyskinesias.

Conclusion: Our patients are younger compared to most series with high prevalence of early onset forms. In the majority of cases, Levodopa was introduced within the first year following onset which expose patients to dyskinesias early in the course of the disease.

Keywords: Parkinson's disease, Clinical features, Treatment, Motor complications, On-motor complications

Introduction

Parkinson's disease (PD) is the second most common neurodegenerative disorder. In general population, its prevalence varies between 18 and 234 cases/100000 people [1].

The mean age of onset is 58 to 62 years and early onset PD, before 40 years, is rare (< 10%) [2]. Symptoms of PD are due to a progressive loss of nigral neurons causing striatal dopaminergic denervation. The advent of levodopa therapy has improved the quality of care but the evolution

is characterized by motor complications in 75-80% of cases after 5 years, they are represented mainly by dyskinesias and motor fluctuations [3,4]. In addition, the health-related quality of life of PD patients is related not only to their motor disability, but also to their non-motor symptoms of depression, sleep disturbance, bladder and sexual dysfunction.

To our knowledge, there are no studies reporting epidemiological and clinical features of PD in Morocco. Our study tries to give an overview of the clinical presentation and management of Idiopathic Parkinson's disease in Moroccan people, from a movement disorders consultation in Ibn Sina University Hospital of Rabat.

* Correspondence: lamiaelachhab@hotmail.com

†Equal contributors

¹Department of Neurology B and Neurogenetics, Hôpital des Spécialités O.N. O, 6444 Rabat, Morocco

⁵University Mohamed V Souissi, Angle avenue Allal El Fassi et Mfadel Cherkaoui, Al Irfane, Rabat, Morocco

Full list of author information is available at the end of the article

Methods

Our series is a study of medical records of 117 patients with diagnosis of Idiopathic Parkinson's disease consecutively seen in our outpatient clinic from January 2006 to January 2011. The patients recruited are representative of Moroccan population as the university hospital of Rabat is the first one in Morocco and patients are mainly from the north-west of the country. The patients were recruited based on the Queen Square Brain Bank Clinical Criteria at clinical assessment [5]. They were examined by the same neurologist. Their clinical records were analyzed, with a form which specifies the demographic characteristics (age, gender, age of onset, duration of the disease), clinical features (laterality of symptoms, predominant feature: Tremulous or akinetic-rigid or mixed, disease severity assessed by UPDRS motor score (Unified Parkinson's Disease Rating Scale) and Hoehn and Yahr score [6], motor complications (fluctuation, dyskinesia, freezing) and non-motor complications (dysautonomic troubles, sleep disorders, depression, cognitive disturbance, hallucination, delirium), treatment regimen (Levodopa treatment, dopaminergic agonist treatment, Levodopa-agonist combination, anticholinergic treatment), mean Levodopa Equivalent Doses (LED) which expresses different doses of anti-parkinsonian medications on a single scale calculated using Deuschl et al.'s method [7], delay of Levodopa introduction and duration of L-dopa treatment.

Statistical analysis

Categorical variables were expressed as effectifs and percentages and continuous variables were expressed as means \pm SD or median (interquartile range). The 95% confidence intervals (CIs) were estimated. To study risk factors, univariate analyses were first performed using simple logistic regression. Variables with $p < 0.20$ in the univariate analysis were tested in the multivariate analysis. Adjusted odds ratios (OR) and their 95% CIs were derived. A p value of 0.05 or less was considered to be statistically significant. Data were analyzed using the statistical software SPSS version 13.0.

Results

Demographic characteristics

Mean age of patients was 64 ± 10 years (range 29–85) with masculine predominance (61.5%) and a sex ratio of M/F = 1.6. Mean age of disease onset was 57.5 ± 11 (range 28–80) years and 13.2% of patients presented an early onset PD with onset before 45 years. Median of disease duration was 5 years (range 3–9).

Clinical presentation

Initial symptoms' side onset was right in 56.5% (65) of patients, left in 31.3% (36) and bilateral but asymmetrical in 12.2% (14). The predominant phenotype was tremulous in 40.2% (47) of patients, akineto-rigid in 28.2% (33) and mixed in 31.6% (37). Mean motor UPDRS score was

Table 1 Fluctuations associated factors

Independent variables	n (%)	Univariate analysis			Multivariate analysis		
		OR	CI 95%	p	OR	CI 95%	P
Age of onset							
> 65 years	2 (7,1)	1					
< 45 years	11 (73,3)	35,7	5,6 - 224	<0,001	40	3,9- 410	0,002
45-54 years	11 (40,7)	8,90	1,75 - 45	0,008	5,80	0,92 - 37	0,06
55- 65 years	8 (19,0)	3,05	0,5 - 15,6	0,17	2,10	0,36-13,1	0,39
Gender							
Men	22 (31,0)						
Women	11 (25,0)	0,70	0,31 - 1,73	0,49			
Disease duration*	5 [3;9]	1,10	1,01 - 1,21	0,019	0,90	0,98-1,008	0,63
Treatment							
Dopaminergic agonists	2 (10,5)	1					
Levodopa	7 (17,1)	1,70	0,3 - 9,3	0,5	0,17	0,01- 2,36	0,18
DA + levodopa	24 (50,0)	8,50	1,7 - 40,8	<0,001	2,10	0,63- 7,3	0,22
LED							
≤ 600 mg/day	7 (11,5)						
> 600 mg/day	26 (55,3)	9,55	3,6- 25,3	<0,001	3,94	1,1- 13,9	0,03
L-dopa treatment duration*	5 [3;9]	0,90	0,96- 1,1	0,32			

*median and quartiles ; n (%) : number (percentage) ;OR : odds-ratio; CI: confidence interval; LED : levodopa equivalent dose.

Table 2 Dyskinesias associated factors

Independent variables	n (%)	Univariate analysis			Multivariate analysis		
		OR	CI 95%	p	OR	CI 95%	p
Age of onset							
> 65 years	4 (13,8)	1					
< 45 years	8 (53,3)	7,1	1,65 – 30,8	0,008			
45-54 years	8 (29,6)	2,6	0,68 - 10	0,15			
55- 65 years	11 (26,2)	2,2	0,62 – 7,8	0,21			
Gender							
Men	22 (30,6)						
Women	9 (20,5)	0,58	0,24 - 1,42	0,2			
Disease duration*	5 [3;9]	1,1	1,02 - 1,22	0,013			
Hoehn and Yahr score							
≤ 2	5 (20,0)	1					
> 2	20 (26,7)	1,4	0,49 - 4,39	0,5			
Treatment							
Dopaminergic agonists	2 (10,5)	1			1		
Levodopa	9 (22,0)	0,41	0,08 - 2,15	0,29	0,4	0,15 -1,06	0,06
DA + levodopa	20 (41,7)	2,5	0,99 - 6,47	0,051	0,14	0,01-1,28	0,08
LED							
≤ 600 mg/day	11 (17,5)	1					
> 600 mg/day	20 (42,6)	3,5	1,46- 8,36	0,005			
L-dopa treatment duration*							
≤5 years	60 (59,4)						
>5 years	41 (40,6)	3,27	1,34- 7,95	0,009			
Délai de début de L-dopa							
>3 months	5 (15,6)	1					
≤ 3 months	25 (35,7)	2,9	1,02- 17,0	0,045	3,2	1,07-9,71	0,03

*median and quartiles ; n (%): number (percentage); OR : odds-ratio; CI: confidence interval; LED: levodopa equivalent dose.

20.5+/-14.5 (0-75). Symptom severity, as measured by the UPDRS score, was mild to moderate in 80% (UPDRS <30) and mean Hoehn and Yahr score was 1.77+/-0.7 (1-3).

There were no differences in the clinical presentation related to the age at onset.

Treatment

Forty four per cent (48) of patients were receiving both dopamine agonists and Levodopa whereas 37.7% (41) received only Levodopa and 18.3% (20) received only dopaminergic agonists.

Anticholinergic drugs were used in 22.3% (26) of patients. The delay of levodopa treatment was 11.6 months (0-120) and 68% (74) of patients received L-dopa as first treatment with mean duration of exposition of 5 years (3-9). In 69% of cases, Levodopa was introduced within the first year following onset. The mean Levodopa Equivalent Doses (LED) was 66 ± 446 mg/day (0-2212).

Motor complications

Motor complications were found in 42% (48) of our patients. Motor fluctuations appeared in 28.7% (33) of patients after 3.7 years of disease duration, dyskinesias in 26.7% (31) after 5.5 years of evolution, freezing in 17.2% (20) and falls in 4.3% (5) of patients.

Motor fluctuations

On univariate analysis, age of onset <45 years (p < 0,001; OR = 35,7; IC = [5,6-224]), duration of the disease (p = 0,019; OR = 1,1; IC = [1,01-1,21]), LED >600 mg/j (p < 0,001; OR = 9,55; IC = [3,6-25,3]) and association of levodopa and dopaminergic agonists (p < 0,001; OR = 8,5; IC = [1,7-40,8]) are the factors associated to the occurrence of motor fluctuations. However, on multivariate analysis, only age of onset and LED were associated to fluctuations. Patients with age of onset <45 develop fluctuations 40 times more comparing to an age > 65 (p = 0,002, OR = 40, IC [3,9- 410]) and motor fluctuations were 4 times more

frequent in patients with LED > 600 mg (p = 0.03; IC [1.1-13.9]) (Table 1).

Dyskinesias

On univariate analysis, age of onset <45 years (p = 0.008; OR = 7.1; IC à 95% = [1.65-30.8]), duration of the disease (p = 0.013; OR = 1.1; IC = [1.02-1.22]), LED > 600 mg/j (p = 0.005; OR = 3.5; IC = [1.46-8.36]), exposition to levodopa duration >5 years (p = 0.009; OR = 3.2; IC = [1.34-7.95]) and delay of levodopa treatment ≤ 3 months (p = 0.04; OR = 2.9; IC = [1.02-17]) were the factors associated to occurrence of dyskinesias. However, considering the same treatment regimen, only early introduction of levodopa increases the risk of dyskinesias by 3.2 with p = 0.03 on multivariate analysis using descending method (Table 2).

Freezing

Freezing is correlated to duration of Levodopa therapy (p = 0.008, OR = 1.01, IC = [1.003-1.02]) and LED >600 mg (p = 0.002). On multivariate analysis, only LED >600 mg is highly associated to freezing (p = 0.006; OR = 4.8; IC = [1.5-14.9]) (Table 3).

Non motor complications

Non motor complications consist mostly in dysautonomic troubles in 44% (51) of cases and sleep disorders in 18.1% (21). Depressive symptoms were found in 12% (14) of cases, cognitive disturbance in 11.2% (13), hallucinations in 8.5% (10) and delirium in 2.6% (3) of patients.

There was no association between duration of the disease and dysautonomic troubles (p = 0.9) nor sleep disturbances (p = 0.3). Furthermore, there was no association between hallucinations and LED, medication type, duration of evolution, depression or cognitive troubles.

Gender phenotype

There were no gender differences in clinical characteristics, demographic features and complications, while there is a difference regarding the Hoehn and Yahr score (≤ 2 in 87.8% of women vs. 65% in men, p = 0.01) (Table 4).

Discussion

This study allowed us to identify some specific features of PD in Moroccan patients.

In fact, our patients were younger compared to other series, 57.5 ± 11 years old versus 61.3 ± 10.9 years old for Lopez *et al.* [8] and 67 ± 11.6 years old for Yust-Katz and *al* [9]. This difference may be explained by high prevalence of early onset PD in our series that suppose a specific genetic profile in our population.

The men predominance is similar to other series [10,8,11,12]. These findings could suggest a clinical difference between men and women but it hasn't been investigated in Parkinson's disease series.

Tremulous phenotype is the most frequent without age intercession as reported by Eun Joo Chung *et al.* [2]. However, Wickremaratchi *et al.* [10], reported an increasing of

Table 3 Freezing associated factors

Independent variables	n (%)	Univariate analysis			Multivariate analysis		
		OR	CI 95%	p	OR	CI 95%	p
Age of onset							
> 65 years	4 (14,3)	1					
< 45 years	5 (33,3)	3	0,66- 13,5	0,15			
45-54 years	5 (17,9)	1,3	0,31- 5,4	0,71			
55- 65 years	6 (14,3)	1	0,25- 3,9	1,00			
Gender	-	0,35	0,10- 1,10	0,078			
Disease duration*	5 [3;9]	1,1	1,03- 1,26	0,009	1	0,99-1,01	0,07
Dysautonomic troubles	13 (26,0)	1,5	0,78- 2,90	0,22			
Cognitive troubles	4 (30,8)	2,3	0,65- 8,70	0,18			
L-dopa treatment duration*	5 [3;9]	1,01	1,003- 1,02	0,008	1	1,1-11,8	0,06
LED (mg) **	667 ± 446	1,001	1,000- 1,002	0,02			
≤ 600 mg	5 (7,8)	1					
> 600 mg	15(32,6)	0,17	0,05- 0,50	0,002	4.8	1,56-14,9	0,006
Treatment							
Dopaminergic agonists	0	1					
Levodopa	9 (22,0)	0,92	0,33- 2,50	0,87			
DA + levodopa	11 (23,4)	0		0,71			

*median and quartiles ;**mean and standard deviation; OR: odds-ratio; CI: confidence interval; LED: levodopa equivalent dose.

Table 4 Gender phenotype

Characteristics	Gender		P
	Men	Women	
Age of onset*			0,32
< 45 years	12 (80,0)	3 (20,0)	
45 to 54 years	17 (60,0)	11 (39,3)	
55 to 65 years	27 (64,3)	15 (35,7)	
> 65 years	15 (51,7)	14 (48,3)	
Predominant feature*			0,50
Tremulous	28 (38,9)	19 (42,2)	
Akineto-rigid	23 (31,9)	10 (22,2)	
Mixed	21 (29,2)	16 (35,6)	
Side onset *			0,41
Right	41 (58,6)	24 (53,3)	
Left	19 (27,1)	17 (37,8)	
Bilateral	10 (14,3)	4 (8,9)	
Hohn and Yahr score*			0,01
≤ 2	39 (65,0)	36 (87,8)	
>2	21 (35,0)	5 (12,2)	
UPDRS score **	21,5 ± 16	19,1 ± 12,6	0,44
Disease duration**	6,99 ± 4,73	6,28 ± 4,31	0,42
Motor fluctuations*	22 (31,0)	11 (25,0)	0,49
Dyskinesias*	22 (30,6)	9 (20,5)	0,23
Freezing*	16 (22,2)	4 (9,1)	0,07
Hallucinations*	9 (12,5)	1 (2,2)	0,08
Dysautonomic troubles*	35 (48,6)	16 (36,4)	0,21
Depression*	9 (12,5)	5 (11,4)	0,85
Cognitive troubles*	8 (11,1)	5 (11,4)	0,96

*number (percentage); **mean ± standard deviation.

tremulous phenotypes with age. In fact, studies are discordant concerning the influence of age on the clinical subtype.

The predominance of symptoms on the right side has been noted in our patients and in other series [9]. Schlomit et al. have reported in their series of 307 patients the asymmetry of symptoms and their predominance on the dominant side [9]. Many hypothesis have been proposed to explain this asymmetry such as genetic predisposition, inborn variation of dopaminergic neurons number in one side, environmental, metabolic or toxic factors [13,14].

Most of our patients (69%) received levodopa as initial treatment. The lack of agonists in Morocco, as only piribedil is available, may explain the initial LevoDopa treatment or its early use when piribedil becomes insufficient in controlling symptoms.

Our patients have moderate handicap scores (Hoehn and Yahr ≤2, motor UPDRS < 30). This early use of levodopa with high dose may explain the moderate severity of Hoehn and Yahr score and UPDRS in our patients as they were assessed while taking their medicine. Indeed,

our mean LED is 1.5 times the one pointed out by Politis et al. with the same duration of disease [12].

Motor complications are represented by motor fluctuations and dyskinesias. Motor fluctuations occurred after 3.7 years of evolution. Similar findings were reported by Lopez et al. [8] which could be explained by the use of levodopa as first line treatment in 69% of patients. However, duration of levodopa therapy is not a risk factor for motor fluctuations in our study while early onset and LED > 600 mg/j were correlated to fluctuations on multivariate analysis.

In other studies, duration of levodopa therapy and severity of the disease (Hoehn and Yahr ≥2) were recognised as risk factors for fluctuations [15].

Concerning dyskinesias, after five years of levodopa therapy, less than one third of our patients presented dyskinesias. This has also been reported in other series in which the prevalence of dyskinesias is 23% [10] while therapeutic trials data suggest that 40% of patients are at risk of developing dyskinesias after 4 to 6 years of levodopa treatment [16]. This difference could be due to the age of patients in therapeutic trials, the treatment regimen and the scores used to identify dyskinesias. In our series, only inaugural use of Levodopa was a risk factor of dyskinesias as it multiplied the risk by 3. On the other hand, early age of onset, LED >600 mg/j and the duration of disease, typically reported in literature have not been identified as risk factors on multivariate analysis [10].

Freezing was rarely found in our patients (17.2%) comparing to the literature. In ancient publications, prevalence of freezing reached 50% after 5 years of evolution [17,18], while in recent series it varies between 26 to 32% [19,20]. This decrease in the prevalence of freezing is most likely due to the use of dopaminergic agonists that allows lowering the doses of levodopa. In fact, the doses of levodopa and severity of the disease are the risk factors of the occurrence of freezing [19,20] and it was also confirmed in our series as a LED > 600 mg/j was the only risk factor.

Non motor complications are represented mainly by dysautonomic troubles (44%) and weren't correlated to disease duration which has also been reported by Sandyk and Awerbuch (1992) [21]. Depression was found only in 12% of cases, while it has been reported in 40 to 70% in other series [22,23]. This difference is probably due to an underestimation of depression in our patients especially that no depression scale was used in our study but the special pathology perception in our population can be another explanation as faith could be a factor of a better acceptance of illness in our population.

As quoted in results, women have less severe disease. Miller and al. reported the same findings with more features such as a later age of onset, tremulous phenotype

predominance, much severe dyskinesias and more depression for women [11].

Our results are limited by the retrospective character of our study and non use of specific scales (especially for non motor complications). Additionally, the study was led in one center, even if it's a specialized one. Yet it's the first Moroccan series describing Parkinson disease and surely a starting point and an experience for other local studies.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

RW and LL participated equally to the work. RW participated in the design of the study and the acquisition of data, LL and RL draft the manuscript, RK and RL participated in the acquisition of data, IF and RR performed the statistical analysis and interpretation of data, AE, BA and YM participated in the coordination of data, AR gave the final approval of the manuscript. All authors read and approved the final manuscript.

Author details

¹Department of Neurology B and Neurogenetics, Hôpital des Spécialités O.N. O, 6444 Rabat, Morocco. ²Laboratory of biostatistics, clinical research and epidemiology, Faculty of medicine and pharmacy of Rabat, Rabat, Morocco. ³Centre de recherche en épidémiologie clinique et essais thérapeutiques Faculty of medicine and pharmacy of Rabat, Rabat, Morocco. ⁴Department of Public Health, Faculty of medicine and pharmacy of Rabat, Rabat, Morocco. ⁵University Mohamed V Souissi, Angle avenue Allal El Fassi et Mfadel Cherkaoui, Al Irfane, Rabat, Morocco.

Received: 30 November 2013 Accepted: 27 February 2014

Published: 6 March 2014

References

1. Chrysostome V, Tison F: **Epidemiologie des syndromes parkinsoniens: Maladie de Parkinson, atrophie multi-systématisée et paralysie supra-nucléaire progressive.** *Rev Neurol (Paris)* 2003, **159**(3):343–352.
2. Chung EJ, Ki CS, Lee WY, Kim IS, Kim JY: **Clinical features and gene analysis in Korean patients with early-onset Parkinson disease.** *Arch Neurol* 2006, **63**:1170–1174.
3. Tanner CM: **Epidemiology of Parkinson's Disease.** *Neurol Clin* 1992, **10**:317–329.
4. Van Gerpen JA, Kumar N, Bower JH, Weigand S, Ahlskog JE: **Levodopa-associated dyskinesia risk among Parkinson disease patients in Olmstead County, Minnesota, 1976–1990.** *Arch Neurol* 2006, **63**:205–209.
5. Vercueil L: **les critères de diagnostic clinique de la maladie de Parkinson et des syndromes parkinsoniens : revue de la littérature.** *Rev Neurol (Paris)* 2000, **156**(S2b):14–22.
6. Goetz CG, Fahn S, Martinez-Martin P, Poewe W, Sampaio C, Stebbins GT, Stern MB, Tilley BC, Dodel R, Dubois B, Holloway R, Jankovic J, Kulisevsky J, Lang AE, Lees A, Leurgans S, LeWitt PA, Nyenhuis D, Olanow CW, Rascol O, Schrag A, Teresi JA, Van Hilten JJ, LaPelle N: **Movement Disorder Society-sponsored revision of the Unified Parkinson's Disease Rating Scale (MDS-UPDRS): process, format, and clinimetric testing plan.** *Mov Disord* 2007, **22**(1):41–47.
7. Deuschl G, Schade-Brittinger C, Krack P, Volkmann J, Schafer H, Botzel K, Daniels C, Deuschländer A, Dillmann U, Eisner W, Gruber D, Hamel W, Herzog J, Hilker R, Klebe S, Kloss M, Koy J, Krause M, Kupsch A, Lorenz D, Lorenzl S, Mehdorn HM, Moringlane JR, Oertel W, Pinsker MO, Reichmann H, Reuss A, Schneider GH, Schnitzler A, Steude U, Sturm V, Timmermann L, Tronnier V, Trottenberg T, Wojtecki L, Wolf E, Poewe W, Voges J, German Parkinson Study Group, Neurostimulation Section: **A randomized trial of deep-brain stimulation for Parkinson's disease.** *N Engl J Med* 2006, **355**(9):896–908. Aug 31.
8. Lopez IC, Ruiz PG, del Pozo SVF: **Motor complications in Parkinson's Disease: ten year follow-up study.** *Mov Disord* 2010, **25**(16):2735–2739.
9. Yust-Katz S, Tesler D, Treves TA, Melamed E, Djaldetti R: **Handedness as a predictor of side of onset of Parkinson's disease.** *Parkinsonism Relat Disord* 2008, **14**:633–635.
10. Wickremaratchi M, Knipe DW, Dwarakanath Sastry BS, Morgan E, Jones A, Salmon R, Weiser R, Moran M, Davies D, Ebenezer L, Raha S, Robertson NP, Butler CC, Ben-Shlomo Y, Morris HR: **The Motor Phenotype of Parkinson's Disease in Relation to Age at Onset.** *Mov Disord* 2011, **26**:457–463.
11. Miller IN, Cronin-Golomb A: **Gender differences in Parkinson's disease: clinical characteristics and cognition.** *Mov Disord* 2010, **25**:2695–2703.
12. Politis M, Wu K, Molloy S, Bain GP, Chaudhuri KR, Piccini P: **Parkinson's disease symptoms: the patient's perspective.** *Mov Disord* 2010, **25**:1646–1651.
13. Djaldetti R, Ziv I, Melamed E: **The mystery of motor asymmetry in Parkinson's disease.** *Lancet Neurol* 2006, **5**(9):796–802.
14. McRitchie DA, Halliday GM, Cartwright H: **Quantitative analysis of the variability of substantia nigra pigmented cell clusters in the human.** *Neuroscience* 1995, **68**:539–551.
15. Hauser RA, McDermott MP, Messing S: **Factors associated with the development of motor fluctuations and dyskinesias in Parkinson disease.** *Arch Neurol* 2006, **63**(12):1756–1760.
16. Ahlskog JE, Muentner MD: **Frequency of levodopa-related dyskinesias and motor fluctuations as estimated from the cumulative literature.** *Mov Disord* 2001, **16**:448–458.
17. Fahn S: **The freezing phenomenon in Parkinsonism.** *Adv Neurol* 1995, **67**:53–63.
18. Giladi N, McMahon D, Przedborski S, Flaster E, Guillery S, Kostic V, Fahn S: **Motor blocks in Parkinson's disease.** *Neurology* 1992, **42**(2):333–339.
19. Giladi N, McDermott MP, Fahn S, Przedborski S, Jankovic J, Stern M, Tanner C, Parkinson Study Group: **Freezing of gait in PD: prospective assessment in the DATATOP cohort.** *Neurology* 2001, **56**:1712–1721.
20. Giladi N, Treves TA, Simon ES, Shabtai H, Orlov Y, Kandinov B, Paleacu D, Kocczyn AD: **Freezing of gait in patients with advanced Parkinson's disease.** *J Neural Transm* 2001, **108**:53–61.
21. Sandyk R, Awerbuch GI: **Dysautonomia in Parkinson's disease: relationship to motor disability.** *Int J Neurosci* 1992, **64**:23–31.
22. Slaughter JR, Slaughter KA, Nichols D, Holmes SE, Martens MP: **Prevalence, clinical manifestations, etiology, and treatment of depression in Parkinson's disease.** *J Neuropsychiatry Clin Neurosci* 2001, **13**:187–196.
23. Leentjens AF, Van den Akker M, Metsemakers JF, Lousberg R, Verhey FR: **Higher incidence of depression preceding the onset of Parkinson's disease: a register study.** *Mov Disord* 2003, **18**:414–418.

doi:10.1186/1755-7682-7-10

Cite this article as: Regragui et al.: Profile of idiopathic parkinson's disease in Moroccan patients. *International Archives of Medicine* 2014 7:10.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

